



MARLETT M 45

Besides being a broad-spectrum contact fungicide, it is also an additional source of Zinc and Manganese for the plants.

Pack Size 100 gm, 250 gm, 500 gm & 1 kg

Specifications

- It is broad spectrum contact fungicide.
- It is having multi-site activity on the cell wall.
- It acts as an additional source on Zn and Mn in the plant.
- Cost effective and gives long duration of control.

Other details

- Category: Dithiocarbamate group of fungicide.
- Mode of action: Multi-site activity.
- Compatibility: It is not compatible with the alkaline chemicals.
- Phytotoxicity: Not phytotoxic when applied as per the recommendations.

Crop Application

Crop	Disease	Dose (gm/acre)
Wheat	Brown & Black Rust, Blight	600-800 gm/acre
Maize	Leaf Blight, Downy Mildew	600-800 gm/acre
Paddy (Rice)	Blast	600-800 gm/acre
Jowar	Leaf Spot	600-800 gm/acre
Potato	Late Blight, Early Blight	600-800 gm/acre
Tomato	Late Blight, Buckeye Rot, Leaf Spot	600-800 gm/acre
Chillies	Damping off	3g (soil drench)
	Fruit Rot, Ripe Rot	600-800 gm/acre
	Leaf Spot	600-800 gm/acre
Cauliflower	Collar Rot	3 gm/acre
	Leaf Spot	600-800 gm/acre
Groundnut	Tikka Disease and Rust	600-800 gm/acre
	Collar Rot, Leaf Spot	25 to 30/10 kg seed
Grapes	Angular Leaf Spot, Downy Mildew, Anthracnose	600-800 gm/acre
Guava	Fruit Rot	20 gm/tree
Banana	Cigar End Rot, Tip Rot, Sigatoka Leafspot	600-800 gm/acre
Apple	Scab & Sooty Blotch	30 gm/tree
Cumin	Blight	600-800 gm/acre

[Sassy_Social_Share]